

$$f(x) = x^3 - x - 4$$

$$f(x_5) = (1.796875)^3 - 1.796875 - 4$$

$$f(x_5) = 3.00 \times 10^{-3} = +ve \quad +ve$$

7th iteration \rightarrow Root lies b/w $x_5 = 1.796875$
and $x_4 = 1.78125$

$$x_6 = \frac{1.796875 + 1.78125}{2}$$

$$x_6 = 1.7890625$$

Put x_6 in $f(x)$

$$f(x) = x^3 - x - 4$$

$$f(x_6) = (1.7890625)^3 - 1.7890625 - 4$$

$$f(x_6) = -0.0672 \quad -ve$$

8th iteration \rightarrow Root lies b/w $x_6 = 1.7890625$
and $x_5 = 1.796875$

$$x_7 = \frac{1.7890625 + 1.796875}{2}$$

$$x_7 = 1.79296$$

Put x_7 in $f(x)$

$$f(1.79296) = -0.02913 \text{ -ve}$$

4th iteration \rightarrow Root lies b/w $x_7 = 1.79296$ & $x_5 = 1.796875$

$$x_8 = \frac{1.79296 + 1.796875}{2}$$

$$x_8 = 1.79491$$

Put x_8 in $f(x)$

$$f(x_8) = -0.02513 \text{ -ve}$$

10th iteration \rightarrow Root lies b/w $x_8 = 1.79491$ & $x_5 = 1.796875$

$$x_9 = \frac{1.79491 + 1.796875}{2}$$

$$x_9 = 1.79589 \checkmark$$

Put x_9 in $f(x)$

$$f(x_9) = -0.00375 \text{ -ve}$$

11th iteration \rightarrow

Root lies b/w $x_9 = 1.79589$ & $x_5 = 1.796875$

$$x_{10} = \frac{1.79589 + 1.796875}{2}$$

$$x_{10} = 1.79638 \checkmark$$

final root 1.796

$$f(x_2) = (1.875)^3 - 1.875 - 4$$

$$f(x_2) = 0.71679 \quad +ve$$

4th iteration \rightarrow Root lies between $x_2 = 1.875$ and $x_1 = 1.75$

$$x_3 = \frac{1.875 + 1.75}{2}$$

$$x_3 = 1.8125$$

Put $x_3 = 1.8125$ in $f(x)$

$$f(x) = x^3 - x - 4$$

$$f(x_3) = (1.8125)^3 - 1.8125 - 4$$

$$f(x_3) = 0.14184 \quad +ve$$

5th iteration \rightarrow Root lies b/w $x_3 = 1.8125$ & $x_1 = 1.75$

$$x_4 = \frac{1.8125 + 1.75}{2}$$

$$x_4 = 1.78125 \quad \checkmark$$

Put x_4 in $f(x)$

$$f(x) = x^3 - x - 4$$

$$f(x_4) = (1.78125)^3 - 1.78125 - 4$$

$$f(x_4) = -0.12960 \quad -ve$$

6th iteration \rightarrow Root lies b/w $x_4 = 1.78125$ & $x_3 = 1.8125$

$$x_5 = \frac{1.78125 + 1.8125}{2}$$

$$x_5 = 1.796875$$

Put x_5 in $f(x)$

* Bisection (OR) Bolzano method \rightarrow

Ques ① \rightarrow find a root of the equation $x^3 - x - 4 = 0$ to four places of decimal by bisection method.

Solⁿ ① \rightarrow $f(x) = x^3 - x - 4 = 0$

$x=0 \rightarrow f(0) = 0 - 0 - 4 = -4 = -ve$ ✓

✓ $x=1 \rightarrow f(1) = 1 - 1 - 4 = -4 = -ve$ ✓

✓ $x=2 \rightarrow f(2) = 8 - 2 - 4 = 8 - 6 = 2 = +ve$ ✓

1st iteration \rightarrow Root lies between $x=1$ & $x=2$

$$x_0 = \frac{1+2}{2} = \frac{3}{2} = 1.5$$

✓ $x_0 = 1.5$

Put $x_0 = 1.5$ in $f(x)$

$$f(x) = x^3 - x - 4$$

$$f(x_0) = (1.5)^3 - 1.5 - 4$$

$$f(x_0) = -2.125 \quad -ve$$

2nd iteration \rightarrow Root lies between $x_0 = 1.5$ & $x = 2$

$$x_1 = \frac{1.5 + 2}{2}$$

$$x_1 = 1.75$$

Put $x_1 = 1.75$ in $f(x)$

$$f(x) = x^3 - x - 4$$

$$f(x_1) = (1.75)^3 - 1.75 - 4$$

$$f(x_1) = -0.390625 \quad -ve$$

3rd iteration \rightarrow Root lies between $x_1 = 1.75$ and $x = 2$

$$x_2 = \frac{1.75 + 2}{2}$$

$$x_2 = \frac{3.75}{2} = 1.875$$

Put x_2 in $f(x)$

$$f(x) = x^3 - x - 4$$

$\therefore f(2.625) = -ve$ and $f(2.75) = +ve$, root lies bet.ⁿ 2.625 & 2.75.

Fourth Approx. $x_4 = \frac{2.625^{(-)} + 2.75^{(+)}}{2} = \underline{\underline{2.6875}}$ Ans
 $f(2.6875) = -0.3391$

$x_5 = 2.71875$

$x_6 = 2.7031$

$x_7 = 2.7109$

$x_8 = 2.707$

$x_9 = 2.7051$

$x_{10} = 2.7061$

$x_{11} = 2.7066$

$x_{12} = 2.7064$

$x_{13} = 2.7065$

$x_{14} = 2.7065$

$\therefore x_{13} = x_{14} = 2.7065$

Hence the root is 2.7065.

Problem 1. Find a root of the equation $x^3 - 4x - 9 = 0$ using the bisection method

(i) in four stages

(ii) correct to four decimal places

Let $f(x) = x^3 - 4x - 9$

$$f(0) = -9$$

$$f(1) = -12$$

$$f(2) = -9$$

$$f(3) = +6$$

$\therefore f(2) = -ve$ and $f(3) = +ve$
so the root lies bet.ⁿ 2 & 3.

First Approx.

$$x_1 = \frac{2+3}{2} = 2.5$$

$$f(2.5) = -3.375$$

~~so~~ $f(2.5) = -ve$ and $f(3) = +ve$,
so the root lies bet.ⁿ 2.5 and 3.

Second Approx.

$$x_2 = \frac{2.5 + 3}{2} = \frac{5.5}{2} = 2.75$$

$$f(2.75) = 0.797 = +ve$$

$\therefore f(2.5) = -ve$ and $f(2.75) = +ve$
so the root lies bet.ⁿ 2.5 & 2.75.

Third Approx.

$$x_3 = \frac{2.5 + 2.75}{2} = 2.625$$

$$f(2.625) = -1.412 = -ve$$

$\therefore f(0.5) = +ve$ and $f(1) = -ve$, the root lies b/w 0.5 & 1

$$x_2 = \frac{0.5 + 1}{2} = 0.75$$

$$f(x_2) = f(0.75) = \cos(57.3 \times 0.75) - (0.75)^{0.75} = -0.8561 = -ve$$

$\therefore f(0.5) = +ve$ & $f(0.75) = -ve$, the root lies b/w 0.5 & 0.75.

$$x_3 = \frac{0.5 + 0.75}{2} = 0.625$$

$$f(x_3) = f(0.625) = -0.3567 = -ve$$

$\therefore f(0.5) = +ve$ & $f(0.625) = -ve$, the root lies b/w
0.5 and 0.625

$$x_4 = \frac{0.5 + 0.625}{2} = 0.5625$$

Hence, the root is 0.5625 Ans

$$f(1) = \cos(57.3) - 1e^1$$

$\therefore f(0) = +ve$ and $f(1) = -ve$, the root lies b/w 0 & 1.

$$x_1 = \frac{0+1}{2} = 0.5;$$

$$f(x_1) = f(0.5) = \cos(57.3 \times 0.5) - (0.5)e^{0.5} = 0.0532 = +ve$$

$\therefore f(0.5) = +ve$ and $f(1) = -ve$, the root lies b/w 0.5 & 1.

$$x_2 = \frac{0.5+1}{2} = 0.75$$

$$f(x_2) = f(0.75) = \cos(57.3 \times 0.75) - (0.75)e^{0.75} = -0.8561 = -ve$$

$\therefore f(0.5) = +ve$ & $f(0.75) = -ve$, the root lies b/w 0.5 & 0.75.

$$x_3 = \frac{0.5+0.75}{2} = 0.625$$

$$f(x_3) = f(0.625) = -0.3567 = -ve$$

$\therefore f(0.5) = +ve$ & $f(0.625) = -ve$, the root lies b/w
0.5 and 0.625

$$x_4 = \frac{0.5+0.625}{2} = 0.5625$$

Q Find a positive root of $\cos x - x e^x = 0$
in four steps.

$$\cos x = x e^x$$

Solⁿ Let $f(x) = \cos x - x e^x$ log₁₀

$$f(0) = \cos(57.3 \times 0) - 0 \cdot e^0 = 1 = +ve$$

$$f(1) = \cos(57.3 \times 1) - 1 \cdot e^1 = -2.18 = -ve$$

$\therefore f(0) = +ve$ and $f(1) = -ve$, the root lies b/w 0 & 1.

$$x_1 = \frac{0+1}{2} = 0.5;$$

$$f(x_1) = f(0.5) = \cos(57.3 \times 0.5) - (0.5)e^{0.5} = 0.0532 = +ve$$

$\therefore f(0.5) = +ve$ and $f(1) = -ve$, the root lies b/w 0.5 & 1.

$$x_2 = \frac{0.5+1}{2} = 0.75$$

$$f(x_2) = f(0.75) = \cos(57.3 \times 0.75) - (0.75)e^{0.75} = -0.8561 = -ve$$

$\therefore f(0.5) = +ve$ & $f(0.75) = -ve$, the root lies b/w 0.5 & 0.75.